

Figure 1. Argentine ants tending mealybugs for honeydew.

Photo: Dong-Hwan Choe, UC Riverside



On Target

A recent survey says pest management professionals working to control ants near structures should consider the micro-environments that are present in the landscape for even better control.

By Herb C. Field, John H. Klotz, Michael K. Rust and Ken Kupfer

Ants are ranked the No. 1 urban pest in the United States by the structural pest control industry, generating an estimated \$1.7 billion annually for pest management professionals (Curl 2005). Most pest management professionals are aware of the difficulties involved in controlling structure-infesting ants: their ability to colonize almost any outdoor environment along with their propensity to invade structures in order to forage or nest makes them particularly challenging.

THE SURVEY. A year-long survey coordinated by the University of California, Riverside and utilizing about 150 pest control technicians from Lloyd Pest Control in San Diego, Calif., was conducted to determine the species of urban pest ants, locations of infestations, and types of control treatments that were applied in residential and commercial settings. The objective was to provide pest management professionals with information that might help them to use their servicing time more wisely and maximize the effectiveness of their treatments.

The survey revealed 18 species of structure-infesting ants, 85 percent of which were Argentine ants (*see Figure 1 above*) (Field et al. 2007). The second most common species was the California harvester ant, which

"It's the first bait I've tried that's both fast and easy to service."

"My service technicians have a huge influence on my final product choices. So I put the Advance[®] Termite Bait System to the test, because my name and business are on the line. With the better station design of Advance, techs appreciate having to spend less time opening difficult stations, replacing broken caps or regaining lost hits when stations are disturbed.

After trying a few homes, I gave my techs the Advance Termite Bait System. Within an hour of sending them out, I was getting calls literally begging me to switch to Advance. I'd never seen such a strong response from the field. Termites take to the bait quicker, stay in the station longer and feed more aggressively than other products."

Mark Drolshagen, Owner
Titan Termite & Pest Control, Charleston, SC



use reader service #55

WHITMIRE MICRO-GEN
Prescription Treatment

Making pest management more practical.

www.advancetbs.com

©2006 Whitmire Micro-Gen

made up 2 percent of the 380 samples that were collected by pest management professionals. In an earlier statewide ant survey of California that was conducted by Knight and Rust (1990), Argentine ants were also the most commonly encountered ant pest by pest management professionals. For example, they found about 40 percent in the San Francisco Bay Area. Undoubtedly Argentine ant numbers and distribution will continue to rise in conjunction with ever-increasing urbanization. Their overwhelming success as an invasive species is due to their ability to produce large supercolonies that through their massive numbers enable them to out-compete other species for limited resources.

Our survey results showed that the majority of the Argentine ant infestations were located in and around residential structures (see Figure 2 below). This may be due to heavy watering and the diversity of plants that homeowners typically grow in southern California. The plants support honeydew-producing insects that flourish in this lush urban environment and provide the Argentine ants with a year-round food supply.

In southern California, full-yard treatments using power sprayers are still a common method to control Argentine ants. However, based on our survey, a better alternative would be sprays directed




Fig. 4. The KM AntPro Sentinel Bait Dispensing System provides 24/7, long-term insect activated low-toxicity bait delivery and monitoring (Rust 2004).

at specific areas around the outside of a structure. Our survey showed that Argentine ants nest and trail primarily along sidewalks and driveways (see Figure 3 on page 78), possibly indicating sources of moisture in these areas — an extremely important resource for ants to survive in the hot, dry summers that are characteristic of this region. Other nesting sites included areas of the lawn adjacent to the foundation, porches, patios, and trees and shrubs. These areas certainly must be part of a treatment program but soil near concrete appears to be critical.

THE TEST. We tested this hypothesis by evaluating several different treatments that were each applied around five homes with heavy infestations of Argentine ants (Klotz et al. 2007). One treatment was liquid bait containing 1 percent disodium octaborate tetrahydrate. We used six to seven KM AntPro bait dispensers (see Figure 4 above) with 20-ounce capacity placed around the outside perimeter of each home and in the yard. After six weeks there was an average of 73 percent reduction of ants tapering off to 58 percent at eight weeks (see Figure 5 on page 80). The increase in ant activity was probably due to ants migrating onto the property from surrounding areas to feed on the bait. This baiting program is designed to target-specific with no residual by taking advantage of the food-sharing behavior of ants to distribute the insecticide throughout the colony. It is a long-term management strategy due to the low concentration of borate being used so the short duration of this study does not adequately reflect its efficacy.

Most effective were the treatments

HOME SWEET HOME



“We switched 7,200 homes in 90 days. Conversion was much easier than we thought.”

“After much investigation and testing, we decided to make the switch from the Sentricon® System to the Advance® Termite Bait System. Although I was prepared to lose some customers, I was delighted to find that we kept 97% of our customers—many of them commenting that they appreciated a free upgrade to better technology. That speaks volumes for Advance and I fully expect this upgrade to increase my retention rate in coming years. I think our customers have noticed the pride our technicians have taken in this improved system. The improved profitability makes me happy. With the support Whitmire Micro-Gen provided to us, I don’t know why any pest control company—large or small—would not convert to ATBS.”

Tony Carder, CEO
Active Pest Control, Atlanta, GA

use reader service #57

WHITMIRE MICRO-GEN
Prescription Treatment
Making pest management more profitable.

www.advancetbs.com

©2006 Whitmire Micro-Gen.

that included Termidor. For example, one treatment consisted of a perimeter spray using 3 to 4 gallons of Termidor (0.06 percent) applied with a backpack sprayer 1 foot up and 1 foot out from the foundation plus along the edges of the sidewalks and driveway. Within one week after treatment, there was a 93 percent reduction in ants, which increased to 98 percent at one month, and then tapered off to 81 percent at two months. Even better control (90 percent reduction at eight weeks) was obtained when this treatment was combined with an application of Talstar granules that was broadcasted into foliage located outside the Termidor zone. But most surprising was the performance of a spot application using only 1 gallon of Termidor applied to active trails of Argentine ants. On average there was a 90 percent reduction of ants eight weeks after treatment.

These results demonstrate the potency of fipronil especially when it is applied to foraging ants, and illustrates the point that ant activity maximizes its effect. This unique compound is non-repellent and sufficiently slow-acting to allow for its pickup


and transfer to other ants. This horizontal transfer of active ingredient is similar to trophallactic exchange of a bait toxicant, except in this case it is due to physical contact with contaminated surfaces, either the substrate the ants are crawling on or another live or dead ant that is contacted (Soeprono and Rust 2004).

CONCLUSIONS. Pest management professionals that are controlling ants near structures should consider the micro-environment

that are present in the landscape. Selection of colony sites by ants is specific to their food and moisture requirements and should be taken into account in the treatment plan. In pest control we can exploit these factors along with the social behavior of ants to our advantage by applying slow-

ANTS WALK THE WALK

Figure 3. The survey showed that Argentine ants nest and trail primarily along sidewalks and driveways.



make a killing



MistAway™ Systems is the leading manufacturer of outdoor misting systems that effectively control mosquitoes, no see ums and other annoying insects.

mosquito control
in a SNAP

We are actively seeking Pest Management Professionals to become MistAway™ dealers.

Contact us today to receive a FREE no obligation informative DVD.

Bug us Anytime

866-485-7255 www.mistaway.com



acting insecticides to the areas frequented by ants so that the active ingredient is picked up and transferred to the rest of the colony.

Herb Field is entomologist and COO, Lloyd Pest Control, San Diego; John Klotz is cooperative extension specialist, University of California-Riverside; Michael Rust is professor of entomology, University of California-Riverside; and Ken Kupfer is president, KM Ant Pro, Nokomis, Fla.

References

Curl, G. 2005. A strategic analysis of the U.S. structural pest control industry – The 2005 season. A survey of Pest Management Professionals in the U.S., Specialty Products Consultants.

Field, H.C., W.E. Evans, R. Hartley, L.D. Hansen, and J.H. Klotz. 2007. A survey of structural ant pests in the southwestern U.S.A. *Sociobiology* 49: 1-14.


Klotz, J.H., M.K. Rust, L. Greenberg, H.C. Field, and K. Kupfer. 2007. An evaluation of several urban pest management strategies to control Argentine ants (Hymenoptera: Formicidae). *Sociobiology* 50: 1-8.

Knight, R.L. and M.K. Rust. 1990. The urban ants of California with distribution notes of imported species. *Southwestern Entomologist* 15: 167-178.

Soeprono, A.M. and M.K. Rust. 2004. Effect of horizontal transfer of barrier insecticides to control Argentine ants (Hymenoptera: Formicidae). *Journal Economic Entomology* 97: 1675-1681.

ANT TREATMENTS OVER 8 WEEKS

Figure 5. Ant numbers are based on the average weight removal (0.3 mg/ant visit) of 50 percent sucrose water from 10 monitoring vials placed around the house for 24 hours. After six weeks there was an average of 73 percent reduction of ants tapering off to 58 percent at eight weeks.



THERE ARE TIMES YOU SHOULD THINK OUTSIDE THE BOX.



NOW IS NOT ONE OF THEM!

MidBox is more than a toolbox. It offers secure and convenient storage, while keeping items out of view and organized. No more hunting for the right tool. Less worry over theft. MidBox provides more cargo and organizational space than a standard toolbox, and is self-contained. It keeps chemicals away from the cab, allowing you to separate and store granules away from liquids. That's thinking safety. MidBox is available for Chevrolet Colorado, GMC Canyon and Ford F-150. For more details see your local dealership.

MIDBOX "INSIDE THE BOX" FEATURES

- Convenient, secure storage
- Seamless appearance
- OEM door handles with matching locks keyed to the vehicle ignition
- Factory direct and available in production colors
- Fully accessorized for your business needs



THINK INSIDE THE BOX



PROFESSIONAL MOBILE STORAGE SOLUTIONS

www.midbox.com / 877-374-5559

use reader service #60